

NOVEL HUMAN TEN-M4/CDZ PROTEINS AND  
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.  
Provisional Application Number 60/275,011, which was filed  
5 on March 12, 2001, and is herein incorporated by reference  
in its entirety.

1. INTRODUCTION

The present invention relates to the discovery,  
identification, and characterization of novel human  
10 polynucleotides encoding proteins that share sequence  
similarity with mammalian membrane proteins. The invention  
encompasses the described polynucleotides, host cell  
expression systems, the encoded proteins, fusion proteins,  
polypeptides and peptides, antibodies to the encoded  
15 proteins and peptides, and genetically engineered animals  
that either lack or overexpress the disclosed  
polynucleotides, antagonists and agonists of the proteins,  
and other compounds that modulate the expression or  
activity of the proteins encoded by the disclosed  
20 polynucleotides, which can be used for diagnosis, drug  
screening, clinical trial monitoring, the treatment of  
diseases and disorders, and cosmetic or nutraceutical  
applications.

2. BACKGROUND OF THE INVENTION

25 Membrane proteins can serve as recognition markers,  
mediate signal transduction, can serve as receptors for  
ligands, and can mediate or facilitate the passage of  
materials across the lipid bilayer. As such, membrane  
proteins are proven drug targets.

30

### 3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with mammalian proteins such as the murine TEN-M4/cdz protein (a dimeric type II transmembrane protein that apparently represents the murine ortholog of the described sequences) and proteins identified as gamma-heregulins.

The novel human nucleic acid sequences described herein encode alternative proteins/open reading frames (ORFs) of 2753 and 1624 amino acids in length (SEQ ID NOS:2 and 4).

The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHPs, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a NHP sequence, or "knock-outs" (which can be conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cell ("ES cell") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique NHP sequences described in SEQ ID NOS:1-4 are

"knocked-out" they provide a method of identifying phenotypic expression of the particular gene, as well as a method of assigning function to previously unknown genes.

In addition, animals in which the unique NHP sequences described in SEQ ID NOS:1-4 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins, which would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody responses. To these ends, gene trapped knockout ES cells have been generated in murine homologs of the described NHPs.

Additionally, the unique NHP sequences described in SEQ ID NOS:1-4 are useful for the identification of protein coding sequences, and mapping a unique gene to a particular chromosome. These sequences identify biologically verified exon splice junctions, as opposed to splice junctions that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology.

Further, the present invention also relates to processes for identifying compounds that modulate, *i.e.*, act as agonists or antagonists of, NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP products, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

30

#### 4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequences of the ORFs encoding the described NHP amino acid sequences, and the encoded NHP amino acid sequences.

5

#### 5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs described for the first time herein are novel proteins that can be expressed in, *inter alia*, human cell lines, human fetal brain, pituitary, spinal cord, lymph node, thyroid, adrenal gland, fetal kidney, fetal lung, and 6- and 9-week old embryos.

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and:

15 (a) nucleotides that encode mammalian homologs of the described nucleotides, including the specifically described NHPs, and the NHP products;

(b) nucleotides that encode one or more portions of the NHPs that correspond to functional domains, and the

20 polypeptide products specified by such nucleotide sequences, including, but not limited to, the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least

25 one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, soluble proteins and peptides in which all or a portion of a hydrophobic transmembrane sequence is deleted; (d) nucleotides that encode chimeric fusion

30 proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e)

therapeutic or diagnostic derivatives of the described polynucleotides, such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs, comprising a sequence first disclosed in the  
5 Sequence Listing.

As discussed above, the present invention includes the human DNA sequences presented in the Sequence Listing (and vectors comprising the same), and additionally contemplates any nucleotide sequence encoding a contiguous  
10 NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and  
15 washing in 0.1x SSC/0.1% SDS at 68°C (Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., N.Y., at p. 2.10.3) and encodes a functionally equivalent expression product. Additionally contemplated are any  
20 nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2x SSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encodes a  
25 functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species, and mutant NHPs, whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as  
30 described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar or identical to corresponding  
5 regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package, as described herein, using standard default settings).

The invention also includes nucleic acid molecules,  
10 preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described herein. In instances where the nucleic acid molecules are  
15 deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80 bases long, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first  
20 disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used  
25 as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a microarray or high-throughput "chip" format). Additionally, a series of NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a  
30 portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS:1-4 can be used as a hybridization probe in conjunction with a

solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (*i.e.*,  
5 gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed  
10 in at least one of the sequences of SEQ ID NOS:1-4, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon, are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637,  
15 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405, the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first  
20 disclosed in SEQ ID NOS:1-4 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of  
25 the production technology. The length of these probes is usually within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides, and more preferably 25 nucleotides, from the sequences first disclosed in SEQ ID NOS:1-4.

30 For example, a series of oligonucleotide sequences, or the complements thereof, can be used in chip format to represent all or a portion of the described NHP sequences. The oligonucleotides, typically between about 16 to about

40 (or any whole number within the stated range)  
nucleotides in length, can partially overlap each other,  
and/or the sequence may be represented using  
oligonucleotides that do not overlap. Accordingly, the  
5 described polynucleotide sequences shall typically comprise  
at least about two or three distinct oligonucleotide  
sequences of at least about 8 nucleotides in length that  
are each first disclosed in the described Sequence Listing.  
Such oligonucleotide sequences can begin at any nucleotide  
10 present within a sequence in the Sequence Listing, and  
proceed in either a sense (5'-to-3') orientation vis-a-vis  
the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of  
broad patterns of genetic activity, providing new  
15 understanding of gene functions, and generating novel and  
unexpected insight into transcriptional processes and  
biological mechanisms. The use of addressable arrays  
comprising sequences first disclosed in SEQ ID NOS:1-4  
provides detailed information about transcriptional changes  
20 involved in a specific pathway, potentially leading to the  
identification of novel components, or gene functions that  
manifest themselves as novel phenotypes.

Probes consisting of sequences first disclosed in SEQ  
ID NOS:1-4 can also be used in the identification,  
25 selection, and validation of novel molecular targets for  
drug discovery. The use of these unique sequences permits  
the direct confirmation of drug targets, and recognition of  
drug dependent changes in gene expression that are  
modulated through pathways distinct from the intended  
30 target of the drug. These unique sequences therefore also  
have utility in defining and monitoring both drug action  
and toxicity.



As an example of utility, the sequences first disclosed in SEQ ID NOS:1-4 can be utilized in microarrays, or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-4 *in silico*, and by comparing previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-4 can be used to identify mutations associated with a particular disease, and also in diagnostic or prognostic assays.

Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be described by the net composition of the nucleotides present within a given region of the sequence, in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in SEQ ID NOS:1-4. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences, can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described

by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent  
5 conditions may refer, e.g., to washing in 6x SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP antisense molecules, useful, for  
10 example, in NHP gene regulation and/or as antisense primers in amplification reactions of NHP nucleic acid sequences. With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple  
15 helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded  
oligonucleotides can additionally comprise at least one modified base moiety that is selected from the group  
20 including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-  
25 galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-  
30 2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine,

5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and  
5 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

10 In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a  
15 phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric oligonucleotide forms specific double-stranded  
20 hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the strands run parallel to each other (Gautier *et al.*, 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.*, 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric  
25 RNA-DNA analogue (Inoue *et al.*, 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g., by use of an  
30 automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein *et al.* (1988, Nucl.

Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), etc.

5 Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, 10 Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (and periodic updates thereof); and Ausubel et al., 1989, *supra*.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using 15 appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single 20 nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification 25 assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

For example, the present sequences can be used in restriction fragment length polymorphism (RFLP) analysis to 30 identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification (as generally described in

U.S. Patent No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e., another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA, prepared from human or non-human cell lines or tissue known to express, or suspected of expressing, an allele of a NHP gene. The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known to express, or suspected of expressing, a NHP gene). A reverse transcription (RT)

reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be  
5 "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can  
10 be used, see e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known  
15 to express, or suspected of expressing, a NHP, in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end  
20 of the normal sequence. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to  
25 that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of  
30 carrying, or known to carry, a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, osteoporosis, obesity, high blood pressure, connective tissue disorders, infertility, etc.), or a cDNA

library can be constructed using RNA from a tissue known to express, or suspected of expressing, a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the  
5 corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

10 Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known to express, or suspected of expressing, a mutant NHP allele in an individual suspected of carrying, or known to carry, such a mutant allele. In this manner, gene products made by the  
15 putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below (for screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A  
20 Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor, N.Y.).

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline  
25 phosphatase fusion proteins. In cases where a NHP mutation results in an expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP expression product.  
30 Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well-known in the art.

The invention also encompasses: (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (*i.e.*, antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculovirus as described in U.S. Patent No. 5,869,336 herein incorporated by reference);

(c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (*i.e.*, gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators, and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include, but are not limited to, the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 or adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP



sequence (transcription factor inhibitors, antisense and ribozyme molecules, or open reading frame sequence or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which  
5 NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists  
10 can be useful for the detection of mutant NHPs, or inappropriately expressed NHPs, for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically  
15 engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells  
20 and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

25 Finally, the NHP products can be used as therapeutics. For example, soluble derivatives, such as NHP peptides/domains corresponding to NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, *i.e.*, fusions of a NHP, or a domain of a NHP, to an IgFc), NHP  
30 antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway), can be used to directly treat diseases,

including infectious diseases, or disorders. For instance, the administration of an effective amount of a soluble NHP, a NHP-IgFc fusion protein, or an anti-idiotypic antibody (or its Fab) that mimics the NHP, could activate or  
5 effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body, delivering a  
10 continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the  
15 invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

## 20 5.1 THE NHP SEQUENCES

The cDNA sequences and the corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. The NHP nucleotides were obtained from clustered genomic sequence, ESTs, and cDNAs produced  
25 using lung and testis mRNAs (Edge Biosystems, Gaithersburg, MD). Sequences corresponding to portions of the presently described NHP sequences have been described, and it is possible that such sequences were incomplete sequences that had been erroneously characterized as "full length."

30 The gene encoding the described NHPs is apparently encoded on one or more of human chromosomes 10 and 11 (see GENBANK accession numbers AC084775 and AP002515). As such,

the described sequences can also be used to map the corresponding coding regions of the human genome.

Several polymorphisms were identified, which include:  
a C/G polymorphism at the region of sequence represented by  
5 nucleotide position 241 of SEQ ID NOS:1 and 3, which can  
result in an arg or gly at corresponding amino acid (aa)  
position 81 of SEQ ID NOS:2 and 4; a C/G polymorphism at  
the region of sequence represented by nucleotide position  
940 of SEQ ID NOS:1 and 3, which can result in a pro or ala  
10 at corresponding aa position 314 of SEQ ID NOS:2 and 4; a  
C/T polymorphism at the region of sequence represented by  
nucleotide position 3465 of SEQ ID NOS:1 and 3, both of  
which result in an asp at corresponding aa position 1155 of  
SEQ ID NOS:2 and 4; and a C/T polymorphism at the region of  
15 sequence represented by nucleotide position 5190 of SEQ ID  
NO:1, both of which result in a gly at corresponding aa  
position 1730 of SEQ ID NO:2. The described NHPs can  
incorporate any and all combinations and permutations of  
the above polymorphisms.

20 An additional application of the described novel  
human polynucleotide sequences is their use in the  
molecular mutagenesis/evolution of proteins that are at  
least partially encoded by the described novel sequences  
using, for example, polynucleotide shuffling or related  
25 methodologies. Such approaches are described in U.S.  
Patent Nos. 5,830,721 and 5,837,458, which are herein  
incorporated by reference in their entirety.

NHP gene products can also be expressed in transgenic  
animals. Animals of any species, including, but not  
30 limited to, worms, mice, rats, rabbits, guinea pigs, pigs,  
micro-pigs, birds, goats, and non-human primates, e.g.,  
baboons, monkeys, and chimpanzees, may be used to generate  
NHP transgenic animals.

Any technique known in the art may be used to introduce a NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (Van der Putten *et al.*, 1985, Proc. Natl. Acad. Sci. USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson *et al.*, 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano *et al.*, 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry a NHP transgene in all their cells, as well as animals that carry a transgene in some, but not all their cells, *i.e.*, mosaic animals or somatic cell transgenic animals. A transgene may be integrated as a single transgene, or in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. A transgene may also be selectively introduced into and activated in a particular cell-type by following, for example, the teaching of Lasko *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous NHP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous NHP gene are

designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous NHP gene (*i.e.*, "knockout" animals).

5       The transgene can also be selectively introduced into a particular cell-type, thus inactivating the endogenous NHP gene in only that cell-type, by following, for example, the teaching of Gu *et al.*, 1994, *Science* 265:103-106. The regulatory sequences required for such a cell-type specific  
10 inactivation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

      Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed  
15 utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may  
20 also be assessed using techniques that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR. Samples of NHP gene-expressing tissue may also be evaluated immunocytochemically using antibodies specific  
25 for the NHP transgene product.

      The present invention also provides for "knock-in" animals. Knock-in animals are those in which a polynucleotide sequence (*i.e.*, a gene or a cDNA) that the animal does not naturally have in its genome is inserted in  
30 such a way that it is expressed. Examples include, but are not limited to, a human gene or cDNA used to replace its murine ortholog in the mouse, a murine cDNA used to replace the murine gene in the mouse, and a human gene or cDNA or

murine cDNA that is tagged with a reporter construct used to replace the murine ortholog or gene in the mouse. Such replacements can occur at the locus of the murine ortholog or gene, or at another specific site. Such knock-in  
5 animals are useful for the *in vivo* study, testing and validation of, *intra alia*, human drug targets, as well as for compounds that are directed at the same and therapeutic proteins.

## 10 5.2 NHPS AND NHP POLYPEPTIDES

NHPS, NHP polypeptides, NHP peptide fragments, mutated, truncated, or deleted forms of the NHPS, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation  
15 of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, and as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical  
20 disorders and diseases. Given the similarity information and expression data, the described NHPS can be targeted (by drugs, oligos, antibodies, etc.) in order to diagnose or treat disease, or to therapeutically supplant or augment the efficacy of, for example, chemotherapeutic agents used  
25 in the treatment of cancer, or agents used in the treatment of inflammatory disorders, arthritis, or infectious disease, or agents used to promote healing.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP sequences. The NHPS  
30 display initiator methionines in DNA sequence contexts consistent with translation initiation sites, but apparently do not display prototypical signal sequences

often associated with other secreted or membrane-associated proteins.

The NHP amino acid sequences of the invention include the amino acid sequence presented in the Sequence Listing, as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention. In fact, any NHP protein encoded by the NHP nucleotide sequences described herein are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing is generically representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al., eds., Scientific American Books, New York, N.Y., herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences, as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation,

etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described herein, but that result  
5 in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example,  
10 nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino  
15 acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the  
20 invention. Where, as in the present instance, the NHP peptide or polypeptide is thought to be membrane protein, the hydrophobic regions of the protein can be excised and the resulting soluble peptide or polypeptide can be recovered from the culture media. Such expression systems  
25 also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such  
30 engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of a NHP, but to assess



biological activity, e.g., in certain drug screening assays.

The expression systems that may be used for purposes of the invention include, but are not limited to,  
5 microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors  
10 containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP nucleotide sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco  
15 mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing NHP nucleotide sequences and  
20 promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors  
25 may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing a NHP, or for raising antibodies to a NHP,  
30 vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to,

the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in-frame with the *lacZ* coding region so that a fusion protein is produced; pIN  
5 vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture Collection) can also be used to  
10 express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa  
15 protease cleavage sites so that the cloned target expression product can be released from the GST moiety.

In an exemplary insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign polynucleotide sequences. The virus grows  
20 in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into a non-essential region (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of a NHP coding sequence  
25 will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence  
30 is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., see Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, may be provided. Furthermore, the initiation codon should be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g.,

5 glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and expression products.

10 Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for the desired processing of the primary transcript, glycosylation, and

15 phosphorylation of the expression product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant

20 proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described herein can be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate

25 expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then

30 switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in turn can be

cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of  
5 compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-  
10 guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes, which can be employed in tk<sup>-</sup>, hgp<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used  
15 as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg,  
20 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147).

25 Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. An exemplary system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991,  
30 Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading

frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni<sup>2+</sup>-nitriloacetic acid-agarose columns, and histidine-  
5 tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct a NHP to a target organ and/or facilitate transport across the membrane into the cytosol.  
10 Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching an appropriate signal sequence to a NHP would also transport a NHP to a desired location within the cell. Alternatively targeting of a NHP  
15 or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes: A Practical Approach", New, R.R.C., ed., Oxford University Press, N.Y., and in U.S. Patent Nos. 4,594,595, 5,459,127, 5,948,767 and  
20 6,110,490 and their respective disclosures, which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of NHPs to a target site or desired organ, where they cross  
25 the cell membrane and/or the nucleus where the NHPs can exert their functional activity. This goal may be achieved by coupling of a NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. Provisional Patent  
30 Application Ser. Nos. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences), to facilitate passage across

cellular membranes, and can optionally be engineered to include nuclear localization signals.

### 5.3 ANTIBODIES TO NHP PRODUCTS

5       Antibodies that specifically recognize one or more epitopes of a NHP, epitopes of conserved variants of a NHP, or peptide fragments of a NHP, are also encompassed by the invention. Such antibodies include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs),  
10 humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

15       The antibodies of the invention may be used, for example, in the detection of a NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of a NHP. Such antibodies may also be  
20 utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction with gene therapy to, for example, evaluate  
25 normal and/or engineered NHP-expressing cells prior to their introduction into a patient. Such antibodies may additionally be used in methods for the inhibition of abnormal NHP activity. Thus, such antibodies may be utilized as a part of treatment methods.

30       For the production of antibodies, various host animals may be immunized by injection with a NHP, a NHP peptide (e.g., one corresponding to a functional domain of a NHP), truncated NHP polypeptides (NHP in which one or

more domains have been deleted), functional equivalents of a NHP or mutated variants of a NHP. Such host animals may include, but are not limited to, pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and/or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin, or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class, including IgG, IgM, IgE, IgA, and IgD, and any subclass thereof. The hybridomas producing the mAbs of this invention may be cultivated *in vitro* or *in*



vivo. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger *et al.*, 1984, Nature, 312:604-608; Takeda *et al.*, 1985, Nature, 314:452-454), by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity, can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patent Nos. 6,114,598, 6,075,181 and 5,877,397 and their respective disclosures, which are herein incorporated by reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies, as described in U.S. Patent No. 6,150,584 and respective disclosures, which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward *et al.*, 1989, Nature 341:544-546) can be adapted to produce single chain antibodies against NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: F(ab')<sub>2</sub>

fragments, which can be produced by pepsin digestion of an antibody molecule; and Fab fragments, which can be generated by reducing the disulfide bridges of F(ab')<sub>2</sub> fragments.

Alternatively, Fab expression libraries may be constructed  
5 (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP,  
10 using techniques well-known to those skilled in the art (see, e.g., Greenspan and Bona, 1993, FASEB J. 7:437-444; and Nissinoff, 1991, J. Immunol. 147:2429-2438). For example, antibodies that bind to a NHP domain and competitively inhibit the binding of NHP to its cognate  
15 receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies, or Fab fragments of such anti-idiotypes, can be used in therapeutic regimens involving a NHP-mediated pathway.

20 Additionally given the high degree of relatedness of mammalian NHPs, the presently described knock-out mice (having never seen a NHP, and thus never been tolerized to a NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the  
25 disclosed mammalian NHPs (i.e., a NHP will be immunogenic in NHP knock-out animals).

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of  
30 the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those

skilled in the art from the foregoing description. Such  
modifications are intended to fall within the scope of the  
appended claims. All cited publications, patents, and  
patent applications are herein incorporated by reference in  
5 their entirety.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230